

(c) *Conditions of use*—(1) *Amount*. 1 bolus (10 grams) daily for a 1,000-pound horse.

(2) *Indications for use*. For treatment of equine diarrhea.

(3) *Limitations*. For horses only; not to be administered to food-producing animals. Do not administer to horses intended for use as food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[48 FR 8054, Feb. 25, 1983, as amended at 50 FR 41489, Oct. 11, 1985]

§ 520.1182 Iron dextran oral suspension.

(a) *Specifications*. Each 1.8 milliliter contains 100 milligrams of elemental iron as ferric hydroxide in complex with a low molecular weight dextran and 0.2 percent phenol as a preservative.

(b) *Sponsor*. See 017800 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount*. 100 milligrams of elemental iron to each pig.

(2) *Indications for use*. Prevention of iron deficiency anemia in baby pigs.

(3) *Limitations*. Treat each pig within 24 hours of farrowing. Administer 1.8 milliliters orally by automatic dose dispenser.

[45 FR 75199, Nov. 14, 1980]

§ 520.1192 Ivermectin paste.

(a) *Specifications*—(1) *Horses*. Paste contains 1.87 percent ivermectin.

(2) *Cattle*. Paste contains 0.153 percent ivermectin.

(b) *Sponsor*. See No. 000006 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Horses*—(i) *Amount*. 200 micrograms per kilogram (91 micrograms per pound) of body weight.

(ii) *Indications for use*. It is used in horses for the treatment and control of large strongyles (adult) (*Strongylus equinus*), (adult and arterial larval stages) (*Strongylus vulgaris*), (adult and migrating tissue stages) (*Strongylus edentatus*), (adult) (*Triodontophorus* spp.); small strongyles, including those resistant to some benzimidazole class compounds (adult and fourth stage larvae) (*Cyathostomum* spp., *Cylicocyclus* spp., *Cylicodontophorus* spp.,

Cylicostephanus spp.); pinworms (adult and fourth stage larvae) (*Oxyuris equi*); ascarids (third- and fourth-stage larvae and adults) (*Parascaris equorum*); hairworms (adult) (*Trichostrongylus axei*); large mouth stomach worms (adult) (*Habronema muscae*); stomach bots (oral and gastric stages) (*Gastrophilus* spp.); lungworms (adults and fourth stage larvae) (*Dictyocaulus arnfieldi*); intestinal threadworms (adults) (*Strongyloides westeri*); summer sores caused by *Habronema* and *Draschia* spp. cutaneous third stage larvae; and dermatitis caused by neck threadworm microfilariae (*Onchocerca* spp.).

(iii) *Limitations*. For oral use only. Do not use in horses intended for food purposes. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(2) *Cattle*—(i) *Amount*. 23 milligrams per 250 pounds of body weight.

(ii) *Indications for use*. It is used in cattle for the treatment and control of gastrointestinal roundworms (adults and fourth-stage larvae) (*Ostertagia ostertagi* (including inhibited forms), *O. lyrata*, *Haemonchus placei*, *Trichostrongylus axei*, *T. colubriformis*, *Cooperia oncophora*, *C. punctata*, *Nematodirus helvetianus*, *Bunostomum phlebotomum*, *Strongyloides papillosus* (adults only), *Oesophagostomum radiatum*, *Trichuris ovis* (adults only)); lungworms (adults and fourth-stage larvae) (*Dictyocaulus viviparus*); grubs (first, second, and third instars) (*Hypoderma bovis*, *H. lineatum*); and sucking lice (*Linognathus vituli*, *Haematopinus eurysternus*).

(iii) *Limitations*. For oral use only. Do not treat cattle within 24 days of slaughter. Because withdrawal time in milk has not been established, do not use in female dairy cattle of breeding age. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[49 FR 22275, May 29, 1984, as amended at 50 FR 27819, July 8, 1985; 51 FR 44449, Dec. 10, 1986; 53 FR 51273, Dec. 21, 1988]

§ 520.1193 Ivermectin tablets and chewables.

(a) *Specifications*—(1) *Dogs*. Each tablet or chewable contains 68, 136, or 272 micrograms of ivermectin.